PROMPT AND AFTERGLOW EMISSION PROPERTIES OF GAMMA-RAY BURSTS WITH SPECTROSCOPICALLY IDENTIFIED SUPERNOVAE

Yuki Kaneko,¹ Enrico Ramirez-Ruiz,^{2,3,4} Jonathan Granot,⁵ Chryssa Kouveliotou,⁶ Stan E. Woosley,⁴

SANDEEP K. PATEL,^{1,6} EVERT ROL,⁷ JEAN J. M. IN 'T ZAND,⁸ ALEXANDER J. VAN DER HORST,⁹

RALPH A. M. J. WIJERS,⁹ AND RICHARD STROM^{9,10}

Received 2006 July 4; accepted 2006 August 7

ABSTRACT

We present a detailed spectral analysis of the prompt and afterglow emission of four nearby long-soft gamma-ray bursts (GRBs 980425, 030329, 031203, and 060218) that were spectroscopically found to be associated with Type Ic supernovae and compare them to the general GRB population. For each event, we investigate the spectral and luminosity evolution and estimate the total energy budget based on broadband observations. The observational inventory for these events has become rich enough to allow estimates of their energy content in relativistic and subrelativistic form. The result is a global portrait of the effects of the physical processes responsible for producing long-soft GRBs. In particular, we find that the values of the energy released in mildly relativistic outflows appears to have a significantly smaller scatter than those found in highly relativistic ejecta. This is consistent with a picture in which the energy released inside the progenitor star is roughly standard, while the fraction of that energy that ends up in highly relativistic ejecta outside the star can vary dramatically between different events.

Subject headings: gamma rays: bursts — supernovae: general

Online material: color figures

1. INTRODUCTION

The discovery in 1998 of a gamma-ray burst (GRB) coincident with a very unusual supernova (SN) of Type Ic (GRB 980425-SN 1998bw, spectroscopically identified) was a turning point in the study of GRBs, offering compelling evidence that long-soft GRBs (Kouveliotou et al. 1993) are indeed associated with the deaths of massive stars (Galama et al. 1998). The large energy release inferred for the supernova suggested that GRBs are potentially associated with a novel class of explosions, having unusual properties in terms of their energy, asymmetry, and relativistic ejecta. More importantly, however, GRB 980425 provided the first hint that GRBs might have an intrinsic broad range of energies: the total energy output in γ -rays (assuming an isotropic energy release) was only $E_{\gamma,\text{iso}} \approx 7 \times 10^{47}$ ergs, some 4 orders of magnitude less energy than that associated with typical GRBs (Bloom et al. 2003). Finally, the fact that GRB 980425/SN 1998bw was located in a nearby galaxy with redshift z = 0.0085 (Tinney et al. 1998; at 35.6 Mpc it remains the closest GRB to the Earth¹¹)

¹ Universities Space Research Association, NSSTC, VP62, 320 Sparkman Drive, Huntsville, AL 35805; yuki.kaneko@nsstc.nasa.gov.

Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540.

³ Chandra Fellow.

⁴ Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064.

⁵ Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, P.O. Box 20450, Mail Stop 29, Stanford, CA 94309.

⁶ NASA MSFC, NSSTC, VP62, 320 Sparkman Drive, Huntsville, AL 35805.
⁷ Department of Physics and Astronomy, University of Leicester, University Road, Leicester LE1 RH7, UK.

⁸ SRON Netherlands Institute for Space Research, Sorbonnelaan 2, 3584 CA, Utrecht, Netherlands; and Astronomical Institute, Utrecht University, P.O. Box 80,000, 3508 TA, Utrecht, Netherlands.

⁹ Astronomical Institute, University of Amsterdam, Kruislaan 403, 1098SJ Amsterdam, Netherlands.

¹⁰ ASTRON, P.O. Box 2, 7990 AA Dwingeloo, Netherlands.

¹¹ Throughout the paper we assume a cosmology with $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.27$, and $\Omega_{\Lambda} = 0.73$.

gave rise to the possibility that such lower energy bursts might be more common than had previously been thought, but harder to detect due to instrumental sensitivity.

Unfortunately, during the elapsed 8 yr, very few SNe have been observed simultaneously with GRBs. To date, three more nearby GRBs have been unambiguously, spectroscopically identified with supernovae, of which two were discovered in 2003 (GRB 030329-SN 2003dh; Stanek et al. 2003; Hjorth et al. 2003, GRB 031203-SN 2003lw; Tagliaferri et al. 2003) and one in 2006 (GRB 060218-SN 2006aj; Pian et al. 2006). Each of these SNe is of the same unusual type as SN 1998bw. Note, however, that there is weaker photometric evidence that many other GRBs are accompanied by SNe, mainly by identification of a late-time "SN bump" in the GRB optical afterglow light curve (e.g., Bloom et al. 1999; Galama et al. 2000; Zeh et al. 2004; Woosley & Bloom 2006). In this paper, we address only the four events with spectroscopically verified SN associations. Among these, only GRB 060218 (interestingly at 143.2 Mpc for z = 0.0335, the closest after GRB 980425; Mirabal et al. 2006; Pian et al. 2006) had γ -ray energetics somewhat comparable to GRB 980425 and could possibly be added to the intrinsically faint GRB sample. In the case of GRB 030329, the total energy release was at the low end of the typical range ($E_{\gamma,iso} \sim 10^{52}$ ergs), much higher than in the other three events. In fact, SN 2003dh was obscured by the extreme optical brightness of the GRB afterglow and was only detected spectroscopically in the GRB optical light curve. Finally, the total energy release of GRB 031203 was intermediate between that of GRB 980425 and regular GRBs. Ramirez-Ruiz et al. (2005) argued that the faint GRB 031203 was a typical powerful GRB viewed at an angle slightly greater than about twice the half-opening angle of the central jet.

In the present study, we consider the energy released during the GRBs, the afterglow, and the SN explosion for these four events in all wave bands, from γ -rays to radio waves, and we also estimate their kinetic energy content. The properties of the prompt and afterglow emission are described in detail in §§ 2 and 3, respectively,



FIG. 1.—Light curves of GRB 980425 detected with the brightest BATSE LAD (20–2000 keV; *top*) and with the *BeppoSAX* WFC (2–27 keV; *bottom*), plotted with 1 s resolution. The four time intervals used in the current analysis are labeled A, B, C, and D. The dotted lines indicate background levels.

and are compared in § 4. The bolometric energy calculation and the evolution of the explosion responsible for their associated SNe are presented in § 5, while § 6 discusses the combined GRB-SN properties and their potential implications. We conclude in § 7 with a brief summary of our primary results and their implications.

2. PROMPT EMISSION

2.1. GRB 980425

GRB 980425 triggered the Large Area Detectors (LADs; 20-2000 keV) of the Burst and Transient Source Experiment (BATSE) on board the Compton Gamma-Ray Observatory, on 1998 April 25 at 21: 49:09 UT (Kippen et al. 1998). The burst was also observed with the Gamma-Ray Burst Monitor (GRBM; 40-700 keV) and one of the two Wide Field Cameras (WFCs; 2-27 keV) on board the BeppoSAX satellite. Figure 1 shows the brightest BATSE LAD and WFC light curves of the burst. Frontera et al. (2000) performed a broadband spectroscopic analysis of the event using the WFC and GRBM data. However, their analysis did not yield sufficiently constrained spectral parameters, due to the much lower sensitivity and poorer energy resolution of GRBM compared to those of the LADs. We performed here a broadband time-resolved spectral analysis of GRB 980425, combining the LAD and the WFC data, to better constrain the spectral parameters and their evolution during the burst.

We used the High Energy Resolution Burst data of the brightest LAD for this event, with 128 channel energy resolution (Kaneko et al. 2006). This data type starts after the BATSE trigger time, which corresponds to the beginning of interval A, as shown in Figure 1. We modeled the LAD background with a low-order polynomial function on spectra taken ~800 s before and ~200 s after the burst episode. The WFC-2 detected the burst at a fairly large off-axis angle of 18°.7 with a light-collecting area of only 10 cm^2 (6% of the on-axis value). However, since the WFC is an imaging telescope with a coded mask, all imaged data can be corrected for background without ambiguity. In addition to the background correction, the data were corrected for the detector dead time, as well as for the variations of the spectral response over the detector area.

 TABLE 1

 Summary of the Spectral Fit Results of GRB 980425

Time Interval	A^{a}	\mathcal{E}_{peak} (keV)	α	$\chi^2/{ m dof}^{ m b}$
A	17.4 ± 3.6	175 ± 13	-0.12 ± 0.22	134.4/141
В	10.7 ± 1.1	133 ± 8	-1.16 ± 0.09	149.9/141
C	4.2 ± 0.9	34 ± 3	-1.54 ± 0.09	158.8/141
D	$2.0^{+5.9}_{-1.3}$	14^{+3}_{-6}	-1.51 ± 0.36	121.8/141

Note.—The data were fitted with a Comptonization model; all uncertainties are 1 $\sigma.$

^a In units of 10^{-3} photons s⁻¹ cm⁻² keV⁻¹.

^b Degree of freedom (dof).



Fig. 2.—Spectral parameter evolution of GRB 980425. The data points correspond to the intervals indicated in Fig. 1. The uncertainties are 1 σ .

To assure good statistics, we binned both the LAD and WFC data into four time intervals (A, B, C, and D in Fig. 1); two intervals each before and after the WFC light-curve peak, similar to the ones used by Frontera et al. (2000). We analyzed the data from both detectors jointly, using the spectral analysis software RMFIT.¹² Since no significant signal was found above 300 keV in all intervals ($\leq 2.2 \sigma$ detection in each interval), we obtained the best fits using the Comptonization photon model (Kaneko et al. 2006),

$$f(\mathcal{E}) = A\left(\frac{\mathcal{E}}{100 \text{ keV}}\right)^{\alpha} \exp\left[-\frac{(2+\alpha)\mathcal{E}}{\mathcal{E}_{\text{peak}}}\right]$$

where A is the amplitude in photons $s^{-1} \text{ cm}^{-2} \text{ keV}^{-1}$, α is the power-law index, and $\mathcal{E}_{\text{peak}}$ is the photon energy (\mathcal{E}), where $\nu F_{\nu} \propto \mathcal{E}^2 f(\mathcal{E})$ peaks. We found no indication that interstellar absorption was required in the fit, and the absorption was therefore not used. Finally, no normalization factor was needed between the data sets of the two detectors.

The spectral fitting results are summarized in Table 1. Figure 2 displays the evolution of \mathcal{E}_{peak} and α ; both parameters clearly

¹² R. S. Mallozzi, R. D. Preece, & M. S. Briggs, "RMFIT, A Lightcurve and Spectral Analysis Tool," ©2006 Robert D. Preece, University of Alabama at Huntsville.



FIG. 3.—HETE-2 FREGATE light curve of GRB 030329 generated using the publicly available Burst Lightcurve data integrated in 64 ms resolution time bins.



Fig. 4.—*INTEGRAL* IBIS/ISGRI light curve of GRB 031203 in 1 s resolution. The four time intervals used in the current analysis are indicated with the dashed lines. The dotted lines show the background levels.

change from hard to soft, in line with the common GRB trend (e.g., Ford et al. 1995; Crider et al. 1997). Moreover, the hardest part of the burst has an $\mathcal{E}_{\text{peak}}$ of 175 ± 13 keV, which is within 1 σ of the peak of the $\mathcal{E}_{\text{peak}}$ distribution for time-resolved BATSE GRB spectra (Kaneko et al. 2006). Furthermore, we found the $\mathcal{E}_{\text{peak}}$ of the duration-integrated spectrum to be 122 ± 17 keV (1 σ uncertainty), placing the event within 2 σ of the peak of the distribution of BATSE GRBs (Kaneko et al. 2006). We find an isotropic-equivalent total energy emitted in 1–10,000 keV of $\mathcal{E}_{\gamma,\text{iso}} = (9.29 \pm 0.35) \times 10^{47}$ ergs, using a luminosity distance of $d_L = 35.6$ Mpc (for z = 0.0085; Tinney et al. 1998).

The total energy fluence values in the X-ray and γ -ray bands are $S_X(2-30 \text{ keV}) = 1.99 \times 10^{-6} \text{ ergs cm}^{-2}$ and $S_{\gamma}(30-400 \text{ keV}) = 3.40 \times 10^{-6} \text{ ergs cm}^{-2}$, respectively. The ratio of the two, log $(S_X/S_{\gamma}) = -0.23$, makes GRB 980425 an X-ray–rich GRB, following the definition of Lamb et al. (2004) and Sakamoto et al. (2005).

2.2. GRB 030329

GRB 030329 was detected with the FREGATE detector (8– 400 keV) on board the *High-Energy Transient Explorer* (*HETE-2*) on 2003 March 29 at 11 : 37:14 UT (Vanderspek et al. 2004; also Fig. 3). A detailed spectral analysis of the event was performed by Vanderspek et al. (2004), who reported an $\mathcal{E}_{\text{peak}} = 70.2 \pm 2.3 \text{ keV}$ for the duration-integrated spectrum. Based on their spectral parameters and flux values, we estimate $E_{\gamma,\text{iso}} = 1.33 \times 10^{52}$ ergs (1–10,000 keV) using $d_L = 790.8$ Mpc (z = 0.1685; Greiner et al. 2003; see also Table 4).

2.3. GRB 031203

GRB 031203 was detected with the IBIS/ISGRI detector (15– 500 keV) on the *International Gamma-Ray Astrophysical Laboratory* (*INTEGRAL*) satellite on 2003 December 3 at 22:01: 28 UT (Gotz et al. 2003). We obtained the IBIS/ISGRI data from the *INTEGRAL* Science Data Center and processed them with OSA version 5.0. Figure 4 displays the light curve of this event, for which we estimated a $T_{90,\gamma}$ duration of 37.0 ± 1.3 s. We extracted a background spectrum at ~300 s prior to the burst and a source spectrum encompassing the $T_{90,\gamma} \pm 10$ s. In addition, we extracted four time-resolved spectra, with durations determined by requiring their signal-to-noise ratios to be above 35, for sufficient statistics. The four intervals are indicated in Figure 4.

We used XSPEC version 12.2.1 for all spectral analyses of this event. The extracted integral source spectrum (17–500 keV, effective exposure time of 45.9 s) is best described by a single power law of index of -1.71 ± 0.08 , consistent with the results

Time Interval (s)	Amplitude ^a	Photon Index	$\chi^2/{ m dof}$
0–2	6.56 ± 2.46	-1.60 ± 0.09	11.2/7
2-4	10.25 ± 4.26	-1.75 ± 0.11	4.9/7
4-8	4.81 ± 1.66	-1.67 ± 0.09	6.2/7
8-50	0.72 ± 0.56	-1.63 ± 0.20	5.0/7

 TABLE 2

 Summary of the Spectral Fit Results of GRB 031203

 Using a Single Power-Law Model

Note.—All uncertainties are 1 σ .

^a In units of photons s⁻¹ cm⁻² keV⁻¹, at 1 keV.

of Sazonov et al. (2004). To constrain the \mathcal{E}_{peak} , Sazonov et al. (2004) fitted the spectrum with the Band function (two power laws smoothly joined at a break energy; Band et al. 1993) with fixed high-energy photon index and derived $\mathcal{E}_{peak} > 190 \text{ keV}$ (90%) confidence level). Employing the same method, we found $\mathcal{E}_{peak} >$ 71 keV (1 σ) and 36 keV (90%), and a low-energy photon index of $\alpha = -1.39 \pm 0.41$. These results are consistent with Sazonov et al. (2004), since the longer integration time used here (45.9 s vs. 22 s) includes the softer tail portion of the event, while the background contribution is practically negligible. Using the single power-law fit parameters, we estimate $S_X/S_{\gamma} = 0.49$ (Table 4), which is in the middle of the range predicted by Sazonov et al. (2004). Therefore, although the INTEGRAL data alone cannot confirm an X-ray-flash (XRF) nature for GRB 031203, as suggested by Watson et al. (2004), we find the event to be clearly X-ray-rich.

The isotropic-equivalent total emitted energy was estimated by extrapolating the single power-law fit between 1 and 10,000 keV in the source rest-frame (z = 0.105; Prochaska et al. 2004), leading to $E_{\gamma,iso} = (1.67^{+0.04}_{-0.10}) \times 10^{50}$ ergs, using $d_L = 472.6$ Mpc. However, since we assumed no spectral break up to 10 MeV, we consider this estimate to be an upper limit of $E_{\gamma,iso}$. All time-resolved spectra were also best fitted with a single power-law model; the fit results are shown in Table 2, and the power-law index evolution is presented in Figure 5. In accord with Sazonov et al. (2004) and as illustrated in Figure 5, we found no significant spectral evolution within the burst, although spectral softening is suggested by the light curve of this event (Fig. 4; see also § 2.5.2).

2.4. GRB 060218

GRB 060218 was detected with the Burst Alert Telescope (BAT; 15-350 keV) on board the Swift satellite on 2006 February 18 at 3:34:30 UT (Cusumano et al. 2006). The burst was exceptionally long, with a $T_{90,\gamma}$ duration of 2100 \pm 100 s (Campana et al. 2006). The available BAT data types were Event data up to ~ 300 s after the trigger time (T_0) and Detector Plane Histogram (DPH) data, thereafter. The latter data type has a variable time resolution; when in survey mode, the nominal resolution is 300 s, but shorter exposures (with a minimum of 64 s) are available immediately after a burst trigger.¹³ After the burst trigger, the satellite slewed immediately to the BAT burst location, and the X-Ray Telescope (XRT; 0.2-10 keV) started observing the burst around $T_0 + 160$ s. The XRT data were in windowed timing (WT) mode from 161 to 2770 s after T_0 . Figure 6 shows the BAT and XRT WT light curves of GRB 060218. Note the much higher photon flux in the X-ray light curve, which clearly indicates the XRF nature of this event. In fact, the burst was de-



Fig. 5.—Single power-law spectral index evolution of GRB 031203. The data points correspond to the intervals indicated in Fig. 4.

tected with BAT by an image trigger, which typically displays no significant signals in the BAT light curve (Cusumano et al. 2006).

We downloaded the publicly available BAT and XRT data and processed them with FTOOLS version 6.0.4. For the present analysis we used the BAT Event data for the first 150 s of the burst and the BAT DPH and XRT WT data for the rest of the burst. This combination of burst trigger and survey mode led to 14 time bins. The time bins are also shown in Figure 6. Before extracting spectra, we rebinned the DPH data with baterebin to the correct energy calibration edges. Then we created the mask-weighting map for the burst location using batmaskwtimg to extract the maskweighted spectra from the rebinned DPH data. Finally, we extracted the spectra using batbinevt for both the Event and DPH data and generated the corresponding detector response matrices with batdrmgen. For the XRT data, we obtained the screened WT event file and extracted spectra for the 13 different time intervals corresponding to the DPH time intervals. We used xselect and an extraction region of 50". For each spectrum, we generated the corresponding auxiliary response files (ARF) with xrtmkarf, using the default empirical ARF file as an input. The spectra were then grouped so that each energy channel contained a minimum of 20 counts, and the latest available response matrix (ver. 007) was used for the analysis. No pileup corrections were necessary for this event.

We analyzed the BAT and XRT data jointly with XSPEC version 12.2.1, except for the first time interval, for which only BAT Event data were available. The BAT and XRT energy ranges used for the analysis were 15–150 keV and 0.6–9 keV, respectively. In the majority of these intervals, an absorbed power law with an exponential high-energy cutoff was the best fit, albeit with a significant low-energy excess. This excess was also identified by Campana et al. (2006) and was fitted with an additional blackbody (BB) component. We have, therefore, included the BB in all spectral fits where the XRT data were included.

The BAT Event data ($T_0 - 8$ to $T_0 + 140$) spectrum is best fitted with a single power law with a high-energy cutoff, with spectral index of -0.87 ± 0.75 and $\mathcal{E}_{\text{peak}}$ of 24.9 \pm 6.0 keV (see also Table 3). The estimated energy fluence during this interval (0.5–150 keV) is (6.38 \pm 4.00) $\times 10^{-7}$ ergs cm⁻². From $T_0 + 140$ to $T_0 + 2734$, we analyzed the BAT-XRT joint spectrum; the best

¹³ BAT Data Products ver. 3.1 by H. Krimm (see http://www.swift.ac.uk/BAT_Data_Products_v3.1.pdf).



Fig. 6.-Light curve of GRB 060218 seen with Swift BAT (15-350 keV; top) and XRT (0.2-10 keV; bottom). The data are binned with 1.6 and 1.0 s resolution, respectively. The 14 time intervals used in our time-resolved analysis are indicated with dotted lines.

TABLE 3 SUMMARY OF THE JOINT BAT-XRT TIME-RESOLVED SPECTRAL FITS FOR GRB 060218

Time Interval (s)	$N_{\rm H} \ (10^{22} \ {\rm cm}^{-2})$	kT (keV)	$rac{\mathcal{E}_{ ext{peak}}}{(ext{keV})}$	Photon Index	χ^2 /dof	Flux $(0.5-150 \text{ keV})^{a}$ $(10^{-9} \text{ ergs s}^{-1} \text{ cm}^{-2})$
			24.9 ± 6.0	-0.87 ± 0.75	52.4/57	6.38 ± 4.00
140-303	0.47 ± 0.08	0.22 ± 0.05	36.1 ± 7.2	-1.39 ± 0.08	238.3/292	$8.95^{+0.43}_{-1.40}$
304-364	0.66 ± 0.14	0.17 ± 0.03	20.8 ± 4.1	-1.49 ± 0.11	165.7/190	$12.84_{-2.87}^{+0.49}$
406-496	0.62 ± 0.11	0.17 ± 0.02	31.5 ± 5.0	-1.39 ± 0.08	236.4/302	$14.93_{-1.85}^{+0.47}$
496-616	0.58 ± 0.02	0.16 ± 0.02	22.3 ± 2.9	-1.33 ± 0.06	323.6/381	$13.99_{-1.12}^{+0.47}$
616–736	0.51 ± 0.07	0.20 ± 0.02	18.9 ± 2.6	-1.21 ± 0.07	395.6/401	$12.59_{-1.02}^{+0.39}$
736-856	0.64 ± 0.08	0.16 ± 0.02	15.2 ± 2.4	-1.41 ± 0.07	397.1/412	$13.40^{+0.37}_{-1.04}$
856–976	0.66 ± 0.08	0.15 ± 0.01	12.0 ± 2.0	-1.45 ± 0.07	390.2/420	$10.03^{+0.39}_{-1.37}$
976-1256	0.61 ± 0.06	0.15 ± 0.01	5.7 ± 0.4	-1.39 ± 0.09	611.7/567	$11.17_{-0.93}^{+0.23}$
1256–1556	0.65 ± 0.06	0.14 ± 0.01	3.6 ± 0.2	-1.30 ± 0.14	571.2/539	$10.26_{-0.97}^{+0.30}$
1557–1857	0.67 ± 0.06	0.13 ± 0.01	2.4 ± 0.4	-1.47 ± 0.03	483.5/481	$8.61^{+0.39}_{-0.95}$
1857–2157	0.60 fixed	0.14 ± 0.003	1.8 ± 0.7	-1.62 ± 0.15	427.8/433	$6.41^{+0.49}_{-0.62}$
2157–2457 [°]	0.60 fixed	0.14 ± 0.004		-2.45 ± 0.03	394.8/388	$6.34_{-0.27}^{+0.42}$
2457–2734 [°]	0.60 fixed	0.13 ± 0.004		-2.54 ± 0.04	327.1/342	$6.15_{-0.52}^{+0.37}$

Notes.—The spectra were fitted to a power law with high-energy cutoff with a blackbody unless otherwise noted. Note that we present \mathcal{E}_{peak} instead of a cutoff energy. The uncertainties are 1 σ .

^a Unabsorbed flux. Uncertainties are associated with the absorbed flux estimated from the fitted parameters. ^b Only BAT Event data were used.

^c Fitted by a power law.



FIG. 7.—Spectral parameter evolution of GRB 060218. The last two points in the photon index plot (*middle*, *squares*) were estimated by fitting a power law with exponential cutoff with the exponential cutoff energy fixed to the last well-determined value (4.7 keV). All the uncertainties are 1 σ .

fit was an absorbed Band function with an additional BB component of $kT = 0.150 \pm 0.004$ keV. The low- and high-energy spectral index and the peak photon energy values were $\alpha = -1.44 \pm 0.06$, $\beta = -2.54 \pm 0.07$, and $\mathcal{E}_{\text{peak}} = 4.67 \pm 1.15$ keV, respectively. The absorption model used was wabs, with the best-fit value of $N_{\text{H}} = (0.60 \pm 0.02) \times 10^{22} \text{ cm}^{-2}$. The unabsorbed energy fluence (0.5-150 keV) was $1.72^{+0.18}_{-0.78} \times 10^{-5}$ ergs cm⁻², where the errors were estimated from the absorbed flux. We derived an isotropic-equivalent total emitted energy of $E_{\gamma,\text{iso}} = (4.33^{+0.41}_{-1.74}) \times 10^{49}$ ergs in 1–10,000 keV (source frame), using

 $d_L = 143.2$ Mpc (z = 0.0335; Mirabal et al. 2006; Pian et al. 2006).

The results of the time-resolved spectral fits are listed in detail in Table 3. We fitted an absorbed power law with a high-energy cutoff and a BB component in all BAT-XRT joint spectra but the last two, for which a single power law with BB provided adequate fits. The $N_{\rm H}$ values were found to be constant ($N_{\rm H} \sim 0.6 \times 10^{22}$ cm⁻²) throughout the first 10 BAT-XRT joint time intervals and were fixed to that value for the last three spectra, where their best-fit $N_{\rm H}$ values were higher, resulting in excessive values of unabsorbed flux. Fixing $N_{\rm H}$ did not significantly affect the other parameters in the fits. Figure 7 displays the spectral parameter evolution of the 14 time-resolved spectra. We observed a hard-tosoft spectral evolution in the nonthermal spectra, while the BB temperature (kT) remained constant. On the average, the BB component contributes 0.13% of the source flux throughout the burst prompt-emission duration.

2.5. Comparison

The γ -ray properties of the four events are summarized in Table 4, and Figure 8 shows the evolution of their isotropic-equivalent γ -ray luminosity, $L_{\gamma,iso}$ (in 2–500 keV, source-frame energy). For GRB 060218, the contributions from the thermal component are shown separately from the $L_{\gamma,iso}$ of the nonthermal component. We find that GRB 060218 had γ -ray luminosity similar to GRB 980425; both were a few orders of magnitude fainter than the other two events. However, since GRB 060218 lasted considerably longer than GRB 980425, its $E_{\gamma,iso}$ was ~50 times larger.

We should emphasize here that we measure directly only the energy radiated in the direction of the Earth per second per steradian per logarithmic energy interval by a source at luminosity distance d_L . The apparent bolometric luminosity may be quite different from the *true* bolometric luminosity if the source is not isotropic.

To compare the light curves of these events, we fitted a twosided Gaussian function to the main pulses of the light curves. We found that the time profiles of these events are similar to the

	TABLE	4	
 	D		A

SUMMARY OF BROADBAND PROPERTIES OF THE PROMPT AND AFTERGLOW EMISSION OF THE FOUR GRBs, ALONG WITH THE PROPERTIES OF THEIR ASSOCIATED SNe

Quantity	980425 (1998bw)	030329 (2003dh)	031203 (2003lw)	060218 (2006aj)
<i>z</i>	0.0085	0.1685	0.105	0.0335
$T_{90,\gamma}$ (s)	$34.9\pm3.8^{\mathrm{a}}$	22.9^{b}	37.0 ± 1.3	$2100\pm100^{\rm c}$
$S(X, 2-30 \text{ keV})^{d}$	1.99E-6	6.71E-5	8.46E-7	1.09E-5
$S(\gamma, 30-400 \text{ keV})^{d}$	3.41E-6	1.20E-4	1.74E-6	3.09E-6
$S_{\rm X}/S_{\gamma}$	0.58	0.56	0.49	3.54
$E_{\gamma,iso}$ (ergs)	$(9.29 \pm 0.35) \times 10^{47}$	1.33×10^{52}	$1.67^{+0.04}_{-0.10} imes 10^{50}$	$4.33^{+0.41}_{-1.74} \times 10^{49}$
$\mathcal{E}_{\text{peak}}$ (keV)	122 ± 17	70 ± 2^{b}	>71	4.7 ± 1.2
$T_{90,X}^{r}$ (days)	640.1	2.61	79.8	9.31
$E_{\rm X,iso}$ (ergs)	1.67×10^{48}	7.09×10^{50}	$8.27 imes 10^{48}$	6.15×10^{47}
$T_{90,R}$ (days)	68.6	72.4	81.2	67.2
$E_{R \text{ iso}}$ (ergs)	4.21×10^{44}	5.64×10^{46}	1.41×10^{45}	1.09×10^{43}
$T_{90,\rm SN}$ (days)	53.9	31.4	53.6	30.7
$E_{\rm SN,iso}$ (ergs)	2.31×10^{49}	1.81×10^{49}	3.15×10^{49}	9.24×10^{48}

Notes.—The energetics in γ -ray, X-ray, radio, and optical (for SNe) wavelengths correspond to 1–10,000 keV, 0.3–10 keV, 5–7 GHz, and 3000–24000 Å in the source frame, respectively, unless noted.

^a BATSE current catalog (see http://gammaray.nsstc.nasa.gov/batse/grb/catalog/current).

^b Vanderspek et al. (2004).

^c Campana et al. (2006).

^d In units of ergs cm^{-2} .



FIG. 8.—Evolution of the γ -ray isotropic-equivalent luminosity (2–500 keV, source-frame energy) for all four events, in the source-frame time. For GRB 060218, the luminosity of the thermal component is plotted separately. [See the electronic edition of the Journal for a color version of this figure.]

overall profile found for GRBs (Nemiroff et al. 1994), with the rising part having a half-width half-maximum (HWHM) much smaller (<1/2) than the HWHM of the decaying part.

2.5.1. Duration-integrated Spectra

Figure 9 compares the best-fit duration-integrated spectra (2– 500 keV in source frame) of the four events in νF_{ν} , which shows the energy radiated per logarithmic photon energy interval. We



FIG. 9.— Unabsorbed best-fit duration-integrated spectra (*solid lines*) of the four events, overplotted with the deconvolved data (*gray crosses*) in the source-frame energy. The data are binned for display purposes. The analysis tool for the *HETE-2* data of GRB 030329 was not publicly available, and we only show here the spectral model presented in Vanderspek et al. (2004). See also text in § 2.5.1.



FIG. 10.—The \mathcal{E}_{peak} of the duration-integrated spectra of GRBs 980425, 030329, and 060218. For GRB 031203, we plot a 1 σ lower limit. As a comparison, the \mathcal{E}_{peak} distributions of 251 bright BATSE GRBs (Kaneko et al. 2006) and 37 *HETE-2* GRB/XRFs (Sakamoto et al. 2005) are plotted. Only well-constrained \mathcal{E}_{peak} values are included in the distributions.

plot here the unabsorbed spectral models in each case (*solid lines*), together with the spectrally deconvolved, blueshifted data points shown in gray. The data are binned here for display purposes. No absorption correction was necessary in the combined LAD-WFC fit of GRB 980425. We used the published spectrum for GRB 030329, and we extrapolated our best-fit model for GRB 031203 to the lower energy range shown here. For GRB 060218 the absorption correction was significant, as seen from the comparison of the model to the XRT data below 10 keV.

In Figure 10 we compare the νF_{ν} peak energy, $\mathcal{E}_{\text{peak}}$, of the same duration-integrated spectra, to the $\mathcal{E}_{\text{peak}}$ distributions of 251 bright BATSE GRBs (Kaneko et al. 2006) and 37 *HETE-2* GRB/XRFs (Sakamoto et al. 2005). The $\mathcal{E}_{\text{peak}}$ values of the four events are lower (softer) than the average value of the bright BATSE GRBs, but they all seem to fit well within the *HETE-2* $\mathcal{E}_{\text{peak}}$ distribution. Note that the BATSE $\mathcal{E}_{\text{peak}}$ distribution shown here is derived using the brightest BATSE GRBs, which tend to be spectrally harder than dim GRBs (Mallozzi et al. 1995).

Finally, in Figure 11, we show where these four events fall in the \mathcal{E}_{peak} - $\mathcal{E}_{\gamma,iso}$ plane, in comparison to the empirical correlation (dashed line) found by Amati et al. (2002) and Lloyd-Ronning & Ramirez-Ruiz (2002). GRB 980425 lies farthest from the socalled Amati relation. Although GRB 031203 also lies away from the relation, this deviation could be the result of a viewing angle slightly outside the edge of the jet (Ramirez-Ruiz et al. 2005). In fact, we find a smaller lower limit on \mathcal{E}_{peak} for this event than previously reported, so it is possible that, after correcting for an off-axis viewing angle, it would be consistent with the relation (compare our Fig. 11 to the left panel in Fig. 3 of Ramirez-Ruiz et al. 2005). We also note that for GRBs 980425 and 031203 Ghisellini et al. (2006) have recently suggested two other possible scenarios that may cause such a deviation from the Amati relation. One is that the GRB radiation may have passed through a scattering screen, and the other is that we may have missed softer emission from the GRBs that lasts much longer. In both cases, observed spectra could have much higher $\mathcal{E}_{\text{peak}}$ values than the intrinsic ones. Altogether, the four events do not appear to follow a specific pattern or correlation in the $\mathcal{E}_{\text{peak}}$ - $E_{\gamma,\text{iso}}$ plane.



FIG. 11.—Locations of GRBs 980425, 031203, 030329, and 060218 in the $\mathcal{E}_{\text{peak}-\mathcal{E}_{\gamma,\text{iso}}}$ plane. The dashed line indicates the correlation found by Amati et al. (2002). The events presented in Ghirlanda et al. (2004) are also shown here.

2.5.2. Spectral Evolution

Figure 12 shows the hardness ratio evolution of all four events in their source-frame time. The hardness ratio is defined here as the ratio of the energy flux at 50–500 keV to 2–50 keV in the source-frame energy. The hard-to-soft evolution is seen in all events except GRB 031203. The lack of spectral evolution in GRB 031203 is probably due to the fact that this burst was relatively dim and that the observed energy range was fairly narrow. The burst was only detected below 200 keV (see Fig. 4), and the break energy seemed to always lie around or above this energy; therefore, the flux above this energy may have been over-



FIG. 12.—Hardness ratio evolution of all four events in the source-frame time. The hardness ratio is determined in the source-frame energy for each event. The values for GRB 030329 were estimated from the spectral parameters presented in Vanderspek et al. (2004).



FIG. 13.—Evolution of the $\mathcal{E}_{\text{peak}}$ in GRBs 980425, 030329, and 060218, in the source-frame time. The dotted lines show the best-fit power-law decay, $\mathcal{E}_{\text{peak}} \propto t^{\phi}$, with $\phi = -1.46 \pm 0.12$, -1.17 ± 0.08 , and -1.40 ± 0.06 , respectively. For GRBs 980425 and 060218, the first points are excluded in the power-law fits.

estimated. We also show in Figure 13 the evolution of the \mathcal{E}_{peak} in three events (GRBs 980425, 030329, and 060218) for which the \mathcal{E}_{peak} values were determined from their time-resolved spectra. To characterize the decaying behavior, we fitted a power law $(\mathcal{E}_{peak} \propto t^{\phi})$ to each of the data sets. Since we are only interested in the decaying behavior, the first points of GRBs 980425 and 060218 were excluded from the fits. We found $\phi = -1.46 \pm$ $0.12, -1.17 \pm 0.08$, and -1.40 ± 0.06 , respectively; the bestfit power laws are shown as dotted lines in Figure 13. We note that in the external shock model, for example, if \mathcal{E}_{peak} is identified with the typical synchrotron frequency ν_m , it is expected to scale as $t^{-3/2}$ once most of the energy is transferred to the external medium and the self-similar deceleration phase sets in (Blandford & McKee 1976). This decaying index is actually similar to what we observed here, especially when taking into account that the asymptotic power law of $t^{-3/2}$ is expected to be approached gradually rather than immediately.

3. AFTERGLOW EMISSION

3.1. X-Ray Afterglow

Figure 14 shows the evolution of the prompt and afterglow isotropic-equivalent luminosity in 0.3–10 keV ($L_{X,iso}$) in the source-frame time for all four events. We estimated the prompt X-ray luminosity values from the spectral analysis of the prompt γ -ray emission (§ 2) by extrapolating the best-fit γ -ray spectra to the X-ray energy range for each event. The afterglow X-ray luminosity values were obtained in various ways.

For GRB 030329, we derived the values using the data presented in Tiengo et al. (2003, 2004). In the case of GRB 031203, we analyzed the (now) archival *XMM-Newton* and *Chandra* observations. Our analysis results are consistent with those presented in Watson et al. (2004). For GRB 060218, we analyzed the *Swift* XRT photon-counting (PC) mode data, starting ~6200 s after the burst trigger. We extracted two time-averaged spectra, covering 6191–8529 s and $1.1 \times 10^4 – 1.3 \times 10^5$ s after the burst



FIG. 14.—Evolution of the X-ray (0.3-10 keV, source-frame energy) prompt and afterglow luminosity (isotropic equivalent) in the source-frame time. [See the electronic edition of the Journal for a color version of this figure.]

trigger, respectively, to account for spectral evolution, and fitted each spectrum with a single power law. The spectral parameters were then used to convert the count rates to the luminosity values. The PC data after 1.3×10^5 s were associated with relatively large uncertainties because the count rates were approaching the XRT sensitivity limit, and therefore, we used the fitted parameters of the second spectrum above to estimate the luminosity values for these data points. We also include in Figure 14 the two Chandra observations of GRB 060218 at $\sim 10^6$ s after the burst trigger, which were presented in Soderberg et al. (2006). Finally, we reanalyzed the *BeppoSAX* observations of GRB 980425/SN 1998bw, which were initially presented in Pian et al. (2000). We used only the data from the *BeppoSAX* Medium Energy Concentrator Spectrometers and followed closely the procedure described in Pian et al. (2000). The only potentially significant difference between the current and the analysis of Pian et al. (2000) is that we estimated the background using the same observation, while Pian et al. (2000) used calibration blank-sky observations. As a result, we found slightly lower net count rates (1.6-10.0 keV) during the two observations in 1998 April (as compared to Table 1 in Pian et al. 2000), indicating a slightly flatter decay than was initially reported. In addition, we found that the observation beginning on 1998 April 27 contained 6.1 ks less data than previously reported (total of 14,862 s exposure time). The count rates and exposure times for the remaining observations are consistent with Pian et al. (2000).

To investigate the energy dissipation behavior in the X-ray afterglow, we fitted a natural cubic spline function to the $L_{X,iso}$ history for each individual afterglow and estimated the cumulative emitted energy as a function of time. The evolution of the cumulative X-ray afterglow energy for all four events is shown in Figure 15. The integration time intervals varied from event to event, since the data time coverage was different for each event. We defined the integration time (i.e., "afterglow" time interval) of each event to be from the earliest X-ray observation time ($\sim 10^4$ s) to the time of the latest available data point. In the case of GRB 060218, however, it was difficult to determine the starting point of the afterglow, since the prompt emission was extremely soft and long and the X-ray follow-up started much earlier



FIG. 15.—Cumulative total emitted energy (isotropic equivalent) of the four SN-GRBs (*black lines*) in the range 0.3–10 keV (source-frame energy) as a function of source-frame time. As a comparison, the cumulative energy of the 10 *Swift* GRBs with known redshifts published in Nousek et al. (2006) are also shown here in color.

than in the other three events. It is possible that the steep decay observed before $\sim 10^4$ s (see Fig. 14) belongs to the decaying part of the prompt emission; we, therefore, chose the afterglow starting time for this event to be 1.1×10^4 s, which corresponds to the onset of the break in its $L_{\rm X,iso}$ history.

To compare the cumulative energy evolution to regular GRBs, we also show in Figure 15 the cumulative energy evolution of the 10 Swift GRBs with known redshifts that were presented in Nousek et al. (2006). For these 10 events, we converted the $L_{X,iso}$ values in 2–10 keV presented there to the $L_{X,iso}$ values in the energy range we used for our four events here (i.e., 0.3–10 keV), using the same spectral parameters as in Nousek et al. (2006), and again fitted a spline function to each $L_{X,iso}$ evolution. The start and end times of the integration were the first and the last points of the actual observations. Figure 15 shows that the final $E_{\rm X,iso}$ values for most of the 10 Nousek et al. (2006) GRBs span the range $\sim 10^{50} - 10^{52}$ ergs. Of the four SN-GRB events, only GRB 030329 falls within this range, while the other three events fall between $\sim 10^{48}$ and $\sim 10^{49}$ ergs. We note that there exists a selection effect based on the observed photon flux; an event would be more likely to be detected when it is closer to us than farther, for a given intrinsic luminosity. Therefore, the difference could be partially due to the fact that these four events occurred relatively nearby compared to the 10 Swift events. From the cumulative X-ray energy evolution, we also derived $T_{90,X}$ values for all events, which we define here as the time in the source frame during which 90% of the radiated afterglow energy (0.3–10 keV in the source frame) is accumulated. Table 4 displays the comparison of the X-ray afterglow properties of all four events, along with their γ -ray prompt properties. Again for the purpose of comparison, Figure 16 shows the $E_{\gamma,iso}$, $E_{X,iso}$, and $T_{90,X}$ values of the four SN-GRBs, along with those of the 10 Nousek et al. (2006) GRBs. Here we plot the $E_{\gamma, iso}$ values between 20 and 2000 keV (in the source frame). There is a clear correlation between $E_{\gamma, iso}$ and $E_{X,iso}$ ($E_{\gamma,iso} \propto E_{X,iso}^{1.0\pm0.3}$) and an anticorrelation between $E_{\gamma,iso}$ and $T_{90, X}$ ($E_{\gamma,iso} \propto T_{90, X}^{-1.2\pm0.3}$). We also find $T_{90, X} \propto E_{X,iso}^{-0.7\pm0.2}$, which can be interpreted as follows.



FIG. 16.—Comparison of the four SN-GRBs (*filled circles*) with the 10 *Swift* GRBs presented by Nousek et al. (2006; *squares*). (*a*) The $E_{\gamma, iso}$ - $E_{X, iso}$ plane; (*b*) the $E_{\gamma, iso}$ - $T_{90, X}$ plane. The $E_{\gamma, iso}$ values used here are determined between 20 and 2000 keV in the source frame.

Events that have large isotropic-equivalent energy (both in γ -rays, $E_{\gamma,iso}$, and in the kinetic energy of their afterglow, $E_{k,iso}$) have a large $E_{X,iso}$, indicating a reasonably narrow spread in the efficiency of converting the afterglow kinetic energy into radiation. Moreover, they are typically associated with narrow jets. This means that most of their kinetic energy is in relativistic outflows carried by highly relativistic ejecta (with $\Gamma\sim$ 30–50; Granot & Kumar 2006). Therefore, they have a relatively small $T_{90,X}$, the time at which most of the energy is injected into the afterglow shock, which is of the order of 10^3 s $\leq t_{\text{break},2} \leq 10^4$ s (Nousek et al. 2006). On the other hand, events that have a small E_{kiso} and small $E_{\gamma,iso}$ naturally have a small $E_{X,iso}$. Such events also tend to have most of their relativistic outflow energy residing in mildly relativistic ejecta, rather than in highly relativistic ejecta (Granot & Ramirez-Ruiz 2004; Waxman 2004). As a result, most of the kinetic energy is transferred to the afterglow shock at relatively late times, hence a large $T_{90,X}$. Furthermore, since such events tend to be only mildly collimated, the true total energy in relativistic ($\Gamma \gtrsim 2$) ejecta has a significantly smaller spread than the one of the isotropic-equivalent energy $(E_{\gamma,iso} \text{ or } E_{X,iso})$.

3.2. Radio Afterglow

We performed radio observations of GRB 060218 with the Westerbork Synthesis Radio Telescope (WSRT); these results are listed in Table 5. Additional radio data for this burst and radio data for the three other events were collected from the literature (GRB 980425: Frail et al. 2003; Kulkarni et al. 1998; GRB 030329: Berger et al. 2003; Frail et al. 2005; Taylor et al. 2005; van der Horst et al. 2005; Resmi et al. 2005; GRB 031203:



Fig. 17.—Radio light curves for all four events at 4.8/4.9 GHz. The errors associated with the data are very small for GRB 980425 and 030329.

Soderberg et al. 2004; GRB 060218: Soderberg et al. 2006; Kamble et al. 2006). The radio flux light curves in 4.8/4.9 GHz, at which all four events were observed, are shown in Figure 17.

As with the X-ray afterglow, we fitted the flux light curve of each event (in a wide band, where available) with a natural cubic spline, to estimate the cumulative energy evolution emitted in radio. To minimize the effect of scintillation in the radio light curves, we used fewer nodes than data points in the fitting. This resulted in a smooth fit to the data, which retains the overall evolution of the light curves. The curve is forced to be 0 at very early times (<0.01 days) and late times (>500 days) or to be the inferred flux of the host galaxy in the case of GRB 031203 (Soderberg et al. 2004). The fitted light curves were then used to create a grid in time-frequency space to obtain the flux profile in a wider frequency range (Fig. 18). Using this, we constructed the cumulative radio afterglow energy evolution in 5-7 GHz, shown in Figure 19. In Table 4, we also present the comparison of the radio afterglow properties of the four events, along with their

	TABLE 5				
Log of the Radio	OBSERVATIONS	FOR	GRB	06021	8

Date (2006)	ΔT (days since trigger)	Integration Time (hr)	Frequency (GHz)	Flux (µJy)
Feb 21.452–21.951	3.30-3.80	5.9	1.4	89 ± 108
Feb 28.463-28.908	10.31-10.76	5.2	2.3	-35 ± 70
Feb 21.481–21.927	3.33-3.78	5.2	4.9	257 ± 36
Feb 24.688-24.943	6.54-6.79	5.9	4.9	125 ± 38
Feb 28.433-28.932	10.28 - 10.78	5.8	4.9	106 ± 42
Mar 10.642–10.905	20.49-20.76	6.1	4.9	6 ± 34
Apr 1.345–1.844	42.20-42.70	12.0	4.9	8 ± 24

Notes.—The flux calibration has been performed using the source 3C 286 as standard. The observations on February 21 and February 28 were alternated between 1.4 and 4.9 GHz, and 2.3 and 4.9 GHz, respectively, observing in 40 minute blocks at each frequency.



FIG. 18.—Spline fits to the broadband radio light curves for the four events.

 γ -ray prompt and X-ray afterglow properties. Similar to the quantity $T_{90,X}$ defined above, we define $T_{90,R}$ here as the time in the source frame during which 90% of the radiated afterglow energy is accumulated in 5–7 GHz in the source-frame energy. The integration time of the radio afterglow emission is 1–100 days after the burst trigger time, in the rest frame of the source.

As can be seen in Table 4, the isotropic-equivalent energy that is radiated in the radio ($E_{R,iso}$) is $\sim 3.5-5$ orders of magnitude smaller than that in X-rays, $E_{X,iso}$. This is predominantly due to the fact that νF_{ν} typically peaks closer to the X-rays than to the radio,



FIG. 19.—Cumulative isotropic-equivalent total emitted energy in 5–7 GHz (source-frame energy) as a function of source-frame time.

and it is very flat above its peak, while it falls much faster toward lower energies. Another effect that enhances the difference between $E_{R,iso}$ and $E_{X,iso}$ is that the latter is calculated over a much wider energy range (0.3–10 keV vs. 5–7 GHz). Finally, since these are isotropic-equivalent energies, most of the contribution to $E_{R,iso}$ is from significantly later times than for $E_{X,iso}$, and the collimation of the outflow, together with relativistic beaming effects, could result in a much larger $E_{X,iso}$ than $E_{R,iso}$. We note that for at least two (GRB 908425 and GRB 060218) of the SN-GRBs, the isotropic-equivalent emitted energy at optical wavelengths can be (much) larger than $E_{X,iso}$, because it is typically dominated by the contribution from the SNe (powered by radioactive decay) rather than by the GRB afterglow emission (see also § 5).

4. PROMPT AND AFTERGLOW PROPERTIES

The isotropic-equivalent luminosity of GRB X-ray afterglows scaled to t = 10 hr after the burst in the source frame, $L_{X,iso}(10 \text{ hr})$, can be used as an approximate estimator for the energy in the afterglow shock for the following reasons. First, at 10 hr the X-ray band is typically above the two characteristic synchrotron frequencies, ν_m (of the accelerated electron with minimum energy) and ν_c (of the electron, whose radiative cooling time equals the dynamical time), so that the flux has very weak dependence¹⁴ on ϵ_B and no dependence on the external density, both of which are associated with relatively large uncertainties. Second, at 10 hr the Lorentz factor of the afterglow shock is sufficiently small ($\Gamma \sim 10$) so that a large fraction of the jet is visible (out to an angle of $\sim \Gamma^{-1} \sim 0.1$ rad around the line of sight) and local inhomogeneities on small angular scales are averaged out. Finally, the fact that the ratio of $L_{X,iso}(10 \text{ hr})$ and $E_{\gamma,iso}$ is fairly constant

¹⁴ When synchrotron–self-Compton is taken into account, the dependence on ϵ_B becomes much stronger (Granot et al. 2006).



FIG. 20.—Isotropic-equivalent luminosity of SN-GRB X-ray afterglows scaled to t = 10 hr (source frame, 2–10 keV) after the burst trigger as a function of their isotropic γ -ray energy release (20–2000 keV). The redshift for each event is also shown in color (adopted from Nousek et al. 2006).

for most GRBs suggests that both can serve as a reasonable measure of the isotropic-equivalent energy content of the ejected outflow.

Figure 20 shows $L_{X,iso}(10 \text{ hr})$ in the 2–10 keV rest-frame energy range as a function of their isotropic γ -ray energy release $(E_{\gamma,iso}, \text{ in } 20-2000 \text{ keV})$ for the four events, in comparison to regular GRBs. A linear relation, $L_{X,iso}(10 \text{ hr}) \propto E_{\gamma,iso}$, seems to be broadly consistent with the data, probably suggesting a roughly universal efficiency for converting kinetic energy into gamma rays in the prompt emission for these four events and those of "regular" long GRBs. This "universal" efficiency is also likely to be high (i.e., the remaining kinetic energy is comparable to, or even smaller than, the energy dissipated and radiated in the prompt emission). If this is the case, the well-known efficiency problem for long GRBs also persists for SN-GRB events. Not surprisingly, there is also a correlation between the cumulative X-ray afterglow energy and isotropic γ -ray energy (see Fig. 16 and also § 6).

5. SN PROPERTIES

Pian et al. (2006) presented the bolometric luminosity evolution (in optical and near-infrared wavelengths; 3000-24000 Å) for all the SNe associated with the GRBs considered here, namely, SNe 1998bw, 2003dh, 2003lw, and 2006aj. We fitted a cubic spline to the luminosity evolution of each SN (Fig. 3 of Pian et al. 2006) and estimated the cumulative energy emitted by the SN as a function of source-frame time. To prevent the spline fits to diverge after the last real data points (≤ 40 days), we assumed that the luminosity becomes much smaller ($\ll 10^{41}$ ergs s⁻¹) at a much later time (≫1000 days). The cumulative SN energy history is plotted in Figure 21 for all four SNe. We also estimated $T_{90,SN}$, during which 90% of the SN energy is accumulated in 3000-24000 Å since 1 day after the burst, as well as the total energy emitted in the same energy range, $E_{\rm SN,iso}$. These values are presented in Table 4, along with the burst properties in other wavelengths. It should be noted here that in most cases $E_{SN,iso}$ is monopolized by the SN emission, while the GRB afterglow dominates at all other wavelengths. The integration time for the SN cumulative energy was from 1-100 days after the burst trigger. The cumulative energy of the four SNe spans a much narrower range than the one reported for the entire SN Ic class (Mazzali et al. 2006a). Among these four, SN 2006aj is the least energetic,



FIG. 21.—Cumulative isotropic-equivalent total emitted energy in 3000– 24000 Å (source-frame energy) as a function of source-frame time, for all SNe associated with the GRBs presented in this work.

albeit more energetic than other broad-lined SNe and normal SNe (Pian et al. 2006).

Furthermore, from the radio observations of Type Ib/c SNe presented by Weiler et al. (2002), we note that the peak radio luminosity of SN 1998bw observed at 6 cm (5 GHz) was a few orders of magnitude higher than the other Type Ib/c SNe, although the overall time evolution of this event was comparable to the others.

6. DISCUSSION

The compilation of the radiated energy inventory, presented in Table 4 and Figure 22, offers an overview of the integrated effects



FIG. 22.—Summary of the isotropic-equivalent total emitted energy of the prompt and afterglow emission of the four GRBs, along with the properties of their associated SNe. The energetics in γ -ray, X-ray, radio, and optical (for SNe) wavelengths correspond to 1–10,000 keV, 0.3–10 keV, 5–7 GHz, and 3000–24000 Å in the source frame, respectively. [See the electronic edition of the Journal for a color version of this figure.]

GRB	Z	δ <i>T</i> (s)	F_p^{a}	β	$\mathcal{E}_{ ext{peak}}$ (keV)	\mathcal{E}_{\max}^{b} (keV)	$\hat{\tau}^{c}$	Limit A	Limit B
980425	0.0085	12.6	0.00314	>3.5	143	300	330	1.6	2.4
030329	0.1685	5.27	0.70	2.44	52.5	400	$4.9 imes 10^7$	13	27
031203	0.105	8.16	< 0.002	2, 3	>148	200	$1.0 \times 10^{6}, 1.5 \times 10^{5}$	8.9, 3.7	16, 7.5
060218	0.0335	1981	0.020	2.75	8.8	100	0.36		

Notes.—Following the formalism of Lithwick & Sari (2001). The high-energy photon index β and \mathcal{E}_{peak} values are for the peak spectra of duration δT , where the parameter limits are 1 σ . For $\hat{\tau} < 1$ and $\mathcal{E}_{max} < m_e c^2$ (as is the case for GRB 060218) there is no lower limit on the Lorentz factor γ . The derivation of Limit B assumes that the high-energy power law continues significantly above the observed range for the cases presented in the table. For GRB 031203 there is only a lower limit on \mathcal{E}_{peak} , and β is not measured, so we conservatively take \mathcal{E}_{peak} to be equal to its lower limit and show results for two representative values of β . ^a The flux normalization at \mathcal{E}_{peak} , $F_p = f \mathcal{E}_{peak}^{-\beta}$, in units of photons s⁻¹ cm⁻² keV⁻¹. ^b The maximum photon energy with significant detection.

^c The optical depth for a photon of energy $m_e c^2$ for $\gamma \beta \approx 1$.

of the energy transfers involved in all the physical processes of long GRB evolution, operating on scales ranging from AU to parsec lengths. The compilation also offers a way to assess how well we understand the physics of GRBs, by the degree of consistency among related entries. We present here our choices for the energy transfers responsible for the various entries in the inventory. Many of the arguments in this exercise are updated versions of what is in the literature, but the present contribution is the considerable range of consistency checks, which demonstrates that many of the entries in the inventory are meaningful and believable.

In the following, \S 6.1 repeats the minimal Lorentz factor estimates in Lithwick & Sari (2001) using the parameters inferred for all four SN-GRB events in this work. Section 6.2 compares a simple model for the emitted radiation to the observational constraints on the integrated energy radiated in the afterglow phase. Finally, the energy contents in relativistic and subrelativistic form are estimated in \S 6.3 and 6.4, respectively.

6.1. Constraints on the Lorentz Factor

As is well known, the requirement that GRBs are optically thin to high-energy photons yields a lower limit on the Lorentz factor of the expansion. We estimate here the minimal Lorentz factor of the outflow based on our observational analysis, by using the results of Lithwick & Sari (2001). They derived two limits, A and B. Limit A arises from the requirement that the optical depth to pair production in the source for the photon with the highest observed energy, \mathcal{E}_{max} , should be smaller than unity. Limit B arises from the requirement that the Thompson optical depth of the e^+ , e^- pairs that are produced by the high-energy photons in the source does not exceed unity. The required parameters to estimate these limits are the variability time of the γ -ray light curve, δT (which is related to the radius of emission by $R \sim \Gamma^2 c \delta T$), and the photon flux per unit energy, $dN_{\rm ph}/dA \, dt \, d\mathcal{E}_{\rm ph} = f \, \mathcal{E}_{\rm ph}^{-\beta}$ for $\mathcal{E}_{\rm min} < \mathcal{E}_{\rm min}$ $\mathcal{E}_{ph} < \mathcal{E}_{max}$. In the latter, both the normalization f and the highenergy photon index β are needed. Although the result depends on the exact choice of parameters, representative values are presented in Table 6. For each event, we determined δT to be the full width at half-maximum (FWHM) of a two-sided Gaussian function fitted to the prompt γ -ray light curve, the peak of which aligns with the main peak of the light curve. We then analyzed the "peak" spectrum with the integration time of δT to obtain the high-energy photon index β and the $\mathcal{E}_{\text{peak}}$ values shown here. The minimal values of the Lorentz factor derived in this way are generally smaller for less energetic events (in terms of $E_{\gamma, iso}$ or $E_{X,iso}$). For the event with the smallest $E_{\gamma,iso}$, GRB 980425, we find $\Gamma_{\min} \sim 2.4$ if we adopt our lower limit for the highenergy photon index ($\beta > 3.5, 1 \sigma$), while a larger value of β

would lower Γ_{\min} (consistent with the results of Lithwick & Sari 2001).

6.2. Integrated Radiated Energy

The integrated (isotropic equivalent) radiated energy during the afterglow is given by

$$E_{\rm rad,iso} = \int dt \int d\nu L_{\nu,\rm iso}(t)$$

= $\eta_t t_{\rm peak} L_{\rm iso}(t_{\rm peak}) = \eta_t \eta_\nu t_{\rm peak} \nu_{\rm peak}(t_{\rm peak}) L_{\nu,\rm iso}(t_{\rm peak}), \quad (1)$

where t, ν , and $L_{\nu,iso}$ are measured in the cosmological frame of the GRB, t_{peak} is the time when $tL_{\text{iso}}(t)$ peaks, ν is the frequency where $\nu L_{\nu,iso}(t_{peak})$ peaks, and

$$\eta_t = \frac{\int dt \, L_{\rm iso}(t)}{t_{\rm peak} L_{\rm iso}(t_{\rm peak})},\tag{2}$$

$$\eta_{\nu} = \frac{L_{\rm iso}(t_{\rm peak})}{\nu_{\rm peak}(t_{\rm peak})L_{\nu,\rm iso}(t_{\rm peak})},\tag{3}$$

are factors of order unity here (although we typically expect $\eta_t \eta_\nu \sim 10$). Since in practice $\nu L_{\nu,iso}$ peaks near the X-ray band, we can assume that $L_{X,iso}$ only mildly underestimates ${}^{15}L_{iso}$, and the time t_{peak} , when $tL_{\text{iso}}(t)$ peaks, is usually rather close to the time t_X , when $tL_{X,iso}(t)$ peaks. Therefore, $t_XL_{X,iso}(t_X)$ provides a convenient lower limit for $E_{rad,iso}$ within an order of magnitude, although it is still possible that the X-ray observations might have missed the actual time when most of the energy was radiated, resulting in a significant underestimate. The values of $t_{\rm X}L_{\rm X,iso}(t_{\rm X})$ are provided in Table 7. We find that these values are typically

¹⁵ Even if νF_{ν} peaks below the X-rays, it is very flat above its peak, so a significant fraction of the afterglow energy is still radiated in the X-rays.

TABLE 7
The Source-Frame Time, t_X , of the Peak in the $tL_{X,iso}$ History
AND THE CORRESPONDING $t_X L_{X,ISO}(t_X)$ Values for the Four Events

GRB	Z	$t_{\rm X}$ (s)	$t_{\rm X}L_{\rm X,iso}(t_{\rm X})$ (ergs)	$t_{\rm X} L_{\rm X,iso}(t_{\rm X})/E_{\gamma,iso}$
980425 030329 031203 060218	0.0085 0.1685 0.105 0.0335	$\begin{array}{c} 2.4 \times 10^{7} \\ 1.7 \times 10^{4} \\ 3.5 \times 10^{5} \\ 3.4 \times 10^{4} \end{array}$	$\begin{array}{c} 6.3 \times 10^{47} \\ 5.3 \times 10^{50} \\ 2.1 \times 10^{48} \\ 2.9 \times 10^{47} \end{array}$	0.68 0.040 0.013 0.0068

a factor of $\sim 2-3$ smaller than $E_{X,iso}$, suggesting $\eta_t \sim 2-3$. Although η_t is defined for the bolometric luminosity, rather than for the X-ray luminosity, the values derived here are still fairly representative.

6.3. Energy Inventory: Relativistic Form

Supernova remnants are understood reasonably well, despite continuing uncertainty about the initiating explosion; likewise, we hope to understand the afterglow of GRBs, despite the uncertainties about their trigger mechanism. The simplest hypothesis is that the afterglow is due to a relativistic expanding blast wave.¹⁶ The complex time structure of some bursts suggests that the central trigger may continue (i.e., the central engine may remain active) for up to ~100 s. However, at much later times all trace of the initial time structure would be lost. All that matters then is essentially how much energy has been injected and its distribution in angle and in Lorentz factor, $\epsilon(>\Gamma, \theta)$, where $\epsilon \equiv dE/d\Omega$.

6.3.1. Kinetic Energy Content

An accurate estimate of the kinetic energy in the afterglow shock requires detailed afterglow modeling and good broadband monitoring, which enables one to determine the values of the shock microphysical parameters (the electron and magnetic energy equipartition fractions, ϵ_e and ϵ_B , and the shock-accelerated electron power-law index, p). However, even then it provides only a lower limit for the kinetic energy due to the conventional and highly uncertain assumption that all of the electrons are accelerated to relativistic energies (Eichler & Waxman 2005; Granot et al. 2006). Nevertheless, an approximate lower limit on the isotropic-equivalent kinetic energy in the afterglow shock, $E_{k,iso}$, can be obtained from the isotropic-equivalent X-ray luminosity, $L_{\rm X,iso}$, since the typical efficiency of the X-ray afterglow, $\epsilon_{\rm X} \equiv$ $tL_{X,iso}(t)/E_{k,iso}(t)$, is $\leq 10^{-2}$ (Granot et al. 2006); a rough lower limit on $E_{k,iso}$ is obtained by adopting $\epsilon(t_X) \sim 10^{-2}$, where t_X is defined as in the previous section and the values for the four events are presented in Table 7. For GRBs 980425 and 060218, we estimate $E_{k,iso}(t_X) \approx 6 \times 10^{49} \epsilon_{X,-2}^{-1}$ ergs and $E_{k,iso}(t_X) \approx 3 \times 10^{49} \epsilon_{X,-2}^{-1}$ $10^{49}\epsilon_{\rm X,-2}^{-1}$ ergs, respectively, where $\epsilon_{\rm X,-2} = \epsilon_{\rm X}(t_{\rm X})/10^{-2}$. These rough estimates are similar to the energy inferred from a more detailed analysis of the X-ray and radio observations of this event $(E_{k,\text{iso}} \approx 5 \times 10^{49} \text{ ergs}; \text{Waxman 2004})$. For GRB 031203, we find $E_{k,\text{iso}}(t_{\text{X}}) \approx 2 \times 10^{50} \epsilon_{\text{X},-2}^{-1}$ ergs.

For GRB 030329, we derive $E_{k,iso}(t_X) \approx 5 \times 10^{52} \epsilon_{X,-2}^{-1}$ ergs. This estimate is comparable to that from the broadband spectrum at $t \approx 10$ days ($E_{k,iso} \sim 5 \times 10^{52}$ ergs for a uniform density and $\sim 10^{52}$ ergs for a wind; Granot et al. 2005), assuming negligible lateral expansion of the jet (Gorosabel et al. 2006). For rapid lateral expansion, the inferred value of $E_{k,iso}(10 \text{ days})$ is lower ($\sim 1.6 \times 10^{51}$ – 5×10^{51} ergs; Berger et al. 2003; Granot et al. 2005), but should correspond to a similar $E_{k,iso}$ before the jet break time ($t_j \approx 0.5$ days) for a comparable initial half-opening angle. There were, however, several rebrightening episodes observed in the optical afterglow light curve of GRB 030329, between ~ 1.5 days and 1 week after the GRB (Lipkin et al. 2004), suggesting energy injection into the afterglow shock that increased its energy by a factor of ~ 10 (Granot et al. 2003). This would imply that $E_{k,iso}$ at $t_X \approx 2 \times 10^4$ s was a factor of ~ 10 lower than our rough estimate or that $\epsilon_X(t_X)$ is as high as ~0.1. A possible alternative explanation for the relatively high value of $t_X L_{X,iso}(t_X)$ that does not require a high afterglow efficiency (ϵ_X) comes about if a flare in the X-rays was present around t_X and was not detected in the optical light curve at the similar time. In this case, $L_{X,iso}(t_X)$ would be dominated by late-time activity of the central source, rather than by emission from the external shock (Ramirez-Ruiz et al. 2001; Granot et al. 2003).

6.3.2. Minimal Energy Estimates

Here we derive a simple but rather robust estimate for the minimal combined energy in the magnetic field (E_B) and in the relativistic electrons (E_e) that are responsible for the observed synchrotron emission of flux density $F_{\nu,R}$ at some frequency ν_R , $E_{\min} = \min(E_B + E_e)$. This estimate is applied for the subrelativistic flow, to avoid the effects of relativistic beaming and reduce the uncertainty in the geometry of the emitting region. Since the source is not resolved, the energy is estimated near the time of the nonrelativistic transition, $t_{\rm NR}$, where we have a better estimate of the source size.

We follow standard equipartition arguments (Pacholczyk 1970; Scott & Readhead 1977; Gaensler et al. 2005; Nakar et al. 2005). The minimal energy is obtained close to equipartition, when $E_B/E_e = 3/4$. At such late times ($t \sim t_{\rm NR} \gtrsim 10^2$ days) it is easiest to detect the afterglow emission in the radio, so ν_R would typically be in the radio band. Furthermore, $\nu_m(t_{\rm NR})$ is also usually around the radio band, meaning that the electrons radiating in the radio would carry a reasonable fraction of the total energy in relativistic electrons. Still, the total energy of all electrons would be a factor of $\gtrsim 10$ larger than E_e , which would increase the total minimal energy by a factor of $\gtrsim 6$. Since the kinetic energy is expected to be at least comparable to that in the relativistic electrons and in the magnetic field, the total energy is likely to be at least an order of magnitude larger than E_{min} .

Following Nakar et al. (2005) $E_e = N_e \gamma_e m_e c^2$, where N_e is the number of electrons with a synchrotron frequency $\nu_{\text{syn}} \sim \nu_R$ and, therefore, with a Lorentz factor $\gamma_e \approx [2\pi m_e c(1+z)\nu_R/eB]^{1/2}$. Also, since $F_{\nu,R} \approx N_e P_{\nu,\max}(1+z)/4\pi d_L^2$, where $P_{\nu,\max} \approx P_{\text{syn}}(\gamma_e)/\nu_{\text{syn}}(\gamma_e) \approx \sigma_T m_e c^2 B/3e$, $N_e \approx 12\pi d_L^2 eF_{\nu,R}/(1+z)\sigma_T m_e c^2 B$. Estimating the emitting volume as $V = (4\pi/3)R^3/\eta$ with $\eta = 10\eta_1$ $(\eta^{-1}$ is the fraction of the volume of a sphere with the radius *R* of the emitting material that is actually occupied by the emitting material), while in terms of our observed quantity, t_{NR} , $R(t_{\text{NR}}) = act_{\text{NR}}/(1+z)$, where $a = 2a_{0.3}$ (*a* is the average apparent expansion velocity at t_{NR} in the cosmological frame of the GRB/SN in units of the speed of light), we obtain

$$E_{\min} = 6 \times 10^{49} a_{0.3}^{9/7} \eta_1^{-3/7} (1+z)^{-19/14} d_{L28}^{8/7} \\ \times \left(\frac{F_{\nu,R}}{1 \text{ mJy}}\right)^{4/7} \left(\frac{\nu_R}{5 \text{ GHz}}\right)^{2/7} \left(\frac{t_{\text{NR}}}{100 \text{ days}}\right)^{9/7} \text{ ergs.} \quad (4)$$

The resulting estimates of $E_{\rm min}$ for the four events discussed in this paper are given in Table 8, along with $t_{\rm NR}$ and $F_{\nu,R}$ values. We used $\nu_R = 4.86$ GHz. As a check of reasonableness, we calculate the minimal external density that corresponds to a total energy of $10E_{\rm min}$, $n_{\rm min} = 10E_{\rm min}/(4\pi/3)R_{\rm NR}^3m_pc^2$, where $R_{\rm NR} = R(t_{\rm NR}) = act_{\rm NR}/(1+z)$, in which we use our fiducial values of a = 2 and $\eta = 10$. These values are also presented in Table 8. It is useful to compare $E_{\rm min}$ to other energy estimates. For GRB 980425, the X-ray afterglow observations suggest an energy of $\sim 5 \times 10^{49}$ ergs $\sim 30E_{\rm min}$ in a mildly relativistic roughly spherical component (Waxman 2004). For GRB 030329 the total

¹⁶ For GRB 030329 this picture is supported by direct measurements of the angular size of its radio afterglow image (Taylor et al. 2004, 2005), which show a superluminal apparent expansion velocity that decreases with time, in good agreement with the predictions of afterglow models (Oren et al. 2004; Granot et al. 2005).

TABLE 8 Summary of the Estimated Minimal Combined Energy, E_{min} , for All Four Events, with $\nu_R = 4.86$ GHz

GRB	Ζ	t _{NR} (days)	$F_{\nu,R}$ (mJy)	E _{min} (ergs)	$n_{\rm min}$ (cm ⁻³)
980425	0.0085	200	3	$1.5 imes 10^{48}$	0.0022
030329	0.1685	100	3	1.8×10^{49}	0.32
031203	0.105	100	0.35	$3.1 imes 10^{48}$	0.048
060218	0.0335	7	0.16	$2.1 imes 10^{46}$	0.78

kinetic energy at late times is estimated to be $\sim 3.2 \times 10^{50}$ ergs $\sim 100E_{\rm min}$ (Granot et al. 2005). Therefore, it can be seen that, while the ratio of the total kinetic energy in the relativistic outflow ($\gamma\beta > 1$) around $t_{\rm NR}$ and $E_{\rm min}$ is indeed $\gtrsim 10$, it is typically $\lesssim 100$. Moreover, the fact that these different energy estimates are consistent lends some credibility to these models.

Some cautionary remarks are in order. The above calculation is only sketchy and should be taken as an order-of-magnitude estimate at present. For example, the usual assumption that at $t_{\rm NR}$ the flow is already reasonably well described by the Newtonian spherical Sedov-Taylor self-similar solution, for which $\eta \approx 10$, is probably not a very good approximation. Numerical studies show that there is very little lateral expansion of the GRB jets while the flow is relativistic, and therefore it takes at least several dynamical timescales after $t_{\rm NR}$ for the flow to approach spherical symmetry. Furthermore, since the flow is still mildly relativistic at this stage, there is still nonnegligible relativistic beaming of the radiation toward the observer from the forward jet and away from the observer from the counterjet. Altogether, the flux is somewhat enhanced due to this mild beaming, and the fraction of the total solid angle that is occupied by the flow at $t_{\rm NR}$ is still considerably smaller than unity. Consequently, this introduces an uncertainty of at least a factor of a few in the estimate of E_{\min} ; however, since the total energy is expected to be $\gtrsim 10E_{\min}$, it would still be at least a factor of a few larger than our estimate of E_{\min} . The theoretical uncertainty on the dynamics of the flow should improve with time as more detailed numerical studies become available.

Another important uncertainty is in the determination of the nonrelativistic transition time, $t_{\rm NR}$. The better estimate of $E_{\rm min}$ should become obtainable as more well-sampled afterglow observations are made and the modeling gets more precise, so that one can more carefully estimate both $t_{\rm NR}$ and $F_{\nu,R}(t_{\rm NR})$. For GRB 980425 we related $t_{\rm NR}$ to the time at which the X-ray light curve steepens, which likely corresponds to the deceleration time of the mildly relativistic ejecta. For GRB 030329, $t_{\rm NR}$ was selected to roughly correspond to the time at which the radio light curve flattens and also to be consistent with estimates derived from direct size measurements of the event (Oren et al. 2004; Granot et al. 2005). In the other two cases, only a crude estimate of $t_{\rm NR}$ can be made, as it is unclear whether current observations clearly display a signature of the nonrelativistic transition.

6.4. Energy Inventory: Subrelativistic Form

Despite the wide range in energies in relativistic ejecta and the even wider range in $E_{\gamma,iso}$, the total (nonneutrino) energy of the associated SNs in all four events spans, at most, a factor of 10. Most of this energy is in nonrelativistic ($\gamma\beta < 2$) kinetic energy; the integrated light of the SN is negligible. While not standard candles, the optical luminosities of the four SNe at peak are all much brighter than average Type Ib or Ic SNe (Woosley & Bloom 2006; Pian et al. 2006; Ferrero et al. 2006). Since the brightness of Type I supernovae at peak is given by the instantaneous rate of



FIG. 23.—Comparisons of collimation-corrected total emitted γ -ray energy $(E_{\gamma})(a)$ of the four SN-GRBs (*red*) and other GRBs and (*b*) of the SN energy (E_k) of the four GRB-SNe (*red*) and other SNe of the same type. In $(a), E_{\gamma,iso}$ was used as an upper limit for GRBs with no jet angle constraints.

decay of ⁵⁶Ni, the ⁵⁶Ni masses are thus inferred to be in the range $\sim 0.2-0.7 M_{\odot}$.

To produce this much ⁵⁶Ni in a Wolf-Rayet (W-R) star requires a kinetic energy of at least $\sim 2 \times 10^{51}$ ergs, even in lower mass W-R stars (Ensman & Woosley 1988). In higher mass stars, a still greater energy is required for the light curve to peak within two weeks after the maximum. Large kinetic energies are also inferred from detailed models of the explosion, especially the light curve and the velocity histories of spectral features (see Woosley & Bloom 2006 and references therein). In summary, the supernova kinetic energies in the four well-studied events almost certainly lie in the range $2 \times 10^{51} - 2 \times 10^{52}$ ergs. In fact, the range of typical GRB-SNe may be much smaller, with the brightness at peak varying by no more than 1 mag in all four events and the kinetic energy in at least three of the four events (all but SN 2006aj) within a factor of 2 of 1.5×10^{52} ergs (Woosley & Bloom 2006).

Finally, in Figure 23, we compare the collimation-corrected total emitted energy in γ -ray (E_{γ}) and supernova kinetic energy estimates (E_k) of these four SN-GRB events with regular GRBs and other broad-lined (1998bw-like) SNe without GRBs. For GRB 030329, we used the jet angle estimate of $\theta_j \sim 0.083-0.14$ rad (Gorosabel et al. 2006). For the other three events, the isotropic-equivalent emitted energy, $E_{\gamma,iso}$, was used as an upper limit, as there was no observational evidence of jet breaks for these events. The E_{γ} values for 27 regular GRBs were adopted from Ghirlanda et al. (2004), where the $E_{\gamma,iso}$ was again used as an upper limit for GRBs with no θ_j constraints. All the E_k estimates were taken from the literature (SN 1998bw: Woosley et al. 1999; SN 2003dh: Mazzali et al. 2003; SN 2003lw: Deng et al.

2005; SN 2006aj: Mazzali et al. 2006b; SN 2002ap: Mazzali et al. 2002; SN 2003jd: Mazzali et al. 2005; SN 1997dq and SN 1997dq: Mazzali et al. 2004; SN 2005bf: Tominaga et al. 2005; Folatelli et al. 2006). We note that the explosion energy (i.e., SN E_k) is much larger than the energy released as GRBs (E_{γ}) and spans a much narrower range than E_{γ} .

7. CONCLUDING REMARKS

One of the liveliest debated issues associated with GRBs concerns the total energy released during the burster explosion: Are GRBs standard candles? The GRB community has vacillated between accepting the initial claims that the GRB intrinsic luminosity distribution was very narrow (Horack et al. 1994), discounting all standard candle claims, accepting a standard total GRB energy of $\sim 10^{51}$ ergs (Frail et al. 2001), and diversifying GRBs into "normal" and "subenergetic" classes. The important new development is that we now have significant observational support for the existence of a subenergetic population based on the different amounts of relativistic energy released during the initial explosion. A network of theoretical tests lends credibility to this idea. The existence of a wide range of intrinsic energies that we presented in this work may pose challenges to using GRBs as standard candles-it is also worth stating explicitly that, when viewed together, these four events fall away from the Amati relation.

Our results are consistent with the emerging hypothesis that GRBs and XRFs share a common origin in massive W-R stars. The central engine gives rise to a polar outflow with two components (Woosley & Bloom 2006). One large-angle outflow (the SN), containing most of the energy and mass, is responsible for exploding the star and producing the ⁵⁶Ni to make the SN bright. Only a tiny fraction of the material in this component reaches mildly relativistic velocities, which is more narrowly focused. A second outflow component (the GRB jet) occupies a narrower solid angle and probably contains less energy (which can range from comparable to much less), and most of its energy is in material with relativistic velocities (where the typical Lorentz factor of the material that carries most of the energy in this component can vary significantly among SN-GRBs). After it exits the star, internal shocks within this jet and external shocks with the residual wind material around the star make the GRB or XRF and its afterglow. Apparently, the properties of the broad component are not nearly so diverse as those of the core jet (Ramirez-Ruiz & Madau 2004; Soderberg et al. 2006; Woosley & Bloom 2006).

Aloy, M.-A., Ibáñez, J.-M., Miralles, J.-A., & Urpin, V. 2002, A&A, 396, 693

- Amati, L., et al. 2002, A&A, 390, 81
- Band, D. L., et al. 1993, ApJ, 413, 281
- Berger, E., et al. 2003, Nature, 426, 154
- Blandford, R. D., & McKee, C. F. 1976, Phys. Fluids, 19, 1130
- Bloom, J. S., Frail, D. A., & Kulkarni, S. R. 2003, ApJ, 594, 674
- Bloom, J. S., et al. 1999, Nature, 401, 453
- Campana, S., et al. 2006, Nature, 442, 1008
- Crider, A., et al. 1997, ApJ, 479, L39
- Cusumano, G., Barthelmy, S., Gehrels, N., Hunsberger, S., Immler, S., Marshall, F., Palmer, D., & Sakamoto, T. 2006, GCN Circ. 4775, http://gcn.gsfc.nasa.gov/ gcn/gcn3/4775.gcn3
- Deng, J., et al. 2005, ApJ, 624, 898
- Eichler, D., & Waxman, E. 2005, ApJ, 627, 861
- Ensman, L. M., & Woosley, S. E. 1988, ApJ, 333, 754
- Ferrero, P., et al. 2006, A&A, 457, 857
- Folatelli, G., et al. 2006, ApJ, 641, 1039
- Ford, L. A., et al. 1995, ApJ, 439, 307
- Frail, D. A., et al. 2001, ApJ, 562, L55
- . 2003, AJ, 125, 2299
- -. 2005, ApJ, 619, 994

We have argued, using well-known arguments connected with parameters such as opacity and variability timescales, that these less energetic events do not require a highly relativistic outflow. Our best estimates of Lorentz factors, Γ , for these events are in the range 2–10. Indeed, it is much more difficult to produce a jet with very high Lorentz factor, i.e., a high energy loading per baryon, than with a low Lorentz factor. A jet with low Lorentz factor could result even if a jet of relatively pure energy is produced, since it may be loaded with excess baryons by instabilities at its walls as it passes through the star, or if it does not precisely maintain its orientation (Ramirez-Ruiz et al. 2002; Aloy et al. 2002; Zhang et al. 2004). The above considerations suggest that GRBs made by jets with lower Lorentz factors should be quite common in the universe (Granot & Ramirez-Ruiz 2004).

Continued advances in observations will surely yield unexpected revisions and additions to our understanding of GRBs in connection with SNe. Currently, we are attempting to draw broad conclusions from limited observations of exceedingly complex phenomena. However, the big surprise at the moment is that these SN-GRB events appear to be intrinsically different from and much more frequent (Granot & Ramirez-Ruiz 2004; Guetta et al. 2004; Pian et al. 2006) than luminous GRBs, which have been observed in large numbers out to higher redshifts.

We are very grateful to Scott Barthelmy and Takanori Sakamoto for their help with the Swift BAT data analysis, to Rob Preece and Michael Briggs for their help with the BATSE-WFC joint analysis, and to Ersin Göğüş for helpful discussions. We also thank the WSRT staff, in particular, Tony Foley. This work is supported by IAS and NASA under contracts G05-6056Z (Y. K.) and through a Chandra Postdoctoral Fellowship award PF3-40028 (E. R. R.), by the Department of Energy under contract DE-AC03-76SF00515 (J. G.), and by PPARC (E. R.). S. E. W. acknowledges support from NASA (NNG05GG08G) and the DOE Program for Scientific Discovery through Advanced Computing (SciDAC; DE-FC02-01ER41176). R. A. M. J. W is supported by the Netherlands Foundation for Scientific Research (NWO) through grant 639.043.302. This paper benefited from collaboration through an EU-funded RTN, grant HPRN-CT-2002-00294. The Westerbork Synthesis Radio Telescope is operated by ASTRON (Netherlands Foundation for Research in Astronomy) with support from NWO.

REFERENCES

- Frontera, F., et al. 2000, ApJS, 127, 59
- Gaensler, B. M., et al. 2005, Nature, 434, 1104
- Galama, T. J., et al. 1998, Nature, 395, 670
- 2000, ApJ, 536, 185
- Ghirlanda, G., Ghisellini, G., & Lazzati, D. 2004, ApJ, 616, 331
- Ghisellini, G., Ghirlanda, G., Mereghetti, S., Bosnjak, Z., Tavecchio, F., & Firmani, C. 2006, MNRAS, 372, 1699
- Gorosabel, J., et al. 2006, ApJ, 641, L13
- Gotz, D., Mereghetti, S., Beck, M., Borkowski, J., & Mowlavi, N. 2003, GCN Circ. 2459, http://gcn.gsfc.nasa.gov/gcn/gcn3/2459.gcn3
- Granot, J., Königl, A., & Piran, T. 2006, MNRAS, 370, 1946
- Granot, J., & Kumar, P. 2006, MNRAS, 366, L13
- Granot, J., Nakar, E., & Piran, T. 2003, Nature, 426, 138
- Granot, J., & Ramirez-Ruiz, E. 2004, ApJ, 609, L9
- Granot, J., Ramirez-Ruiz, E., & Loeb, A. 2005, ApJ, 618, 413
- Greiner, J., Peimbert, M., Estaban, C., Kaufer, A., Jaunsen, A., Smoke, J., Klose, S., & Reimer, O. 2003, GCN Circ. 2020, http://gcn.gsfc.nasa.gov/
- gcn/gcn3/2020.gcn3
- Guetta, D., Perna, R., Stella, L., & Vietri, M. 2004, ApJ, 615, L73
- Hjorth, J., et al. 2003, Nature, 423, 847
- Horack, J. M., et al. 1994, ApJ, 426, L5

- Kamble, A., Ishwara-Chandra, C. H., & Bhattacharya, D. 2006, GCN Circ. 4840, http://gcn.gsfc.nasa.gov/gcn/gcn3/4840.gcn3
- Kaneko, Y., Preece, R. D., Briggs, M. S., Paciesas, W. S., Meegan, C. A., & Band, D. L. 2006, ApJS, 166, 298
- Kippen, R. M., et al. 1998, GCN Circ. 67, http://gcn.gsfc.nasa.gov/gcn/gcn3/ 67.gcn3
- Kouveliotou, C., et al. 1993, ApJ, 413, L101
- Kulkarni, S. R., et al. 1998, Nature, 395, 663
- Lamb, D. Q., Donaghy, T. Q., & Graziani, C. 2004, NewA Rev., 48, 459
- Lipkin, Y. M., et al. 2004, ApJ, 606, 381
- Lithwick, Y., & Sari, R. 2001, ApJ, 555, 540
- Lloyd-Ronning, N., & Ramirez-Ruiz, E. 2002, ApJ, 576, 101
- Mallozzi, R. S., et al. 1995, ApJ, 454, 597
- Mazzali, P. A., et al. 2002, ApJ, 572, L61
- 2003, ApJ, 599, L95
- -. 2004, ApJ, 614, 858
- -. 2005, Science, 308, 1284
- 2006a, ApJ, 645, 1323
- . 2006b, Nature, 442, 1018
- Mirabal, N., et al. 2006, ApJ, 643, L99
- Nakar, E., Piran, T., & Sari, R. 2005, ApJ, 635, 516
- Nemiroff, R. J., et al. 1994, ApJ, 423, 432 Nousek, J. A., et al. 2006, ApJ, 642, 389
- Oren, Y., Nakar, E., & Piran, T. 2004, MNRAS, 353, L35
- Pacholczyk, A. G. 1970, Radio Astrophysics (San Francisco: Freeman)
- Pian, E., et al. 2000, ApJ, 536, 778
- 2006, Nature, 442, 1011
- Prochaska, J. X., et al. 2004, ApJ, 611, 200

- Ramirez-Ruiz, E., Celotti, A., & Rees, M. J. 2002, MNRAS, 337, 1349 Ramirez-Ruiz, E., & Madau, E. 2004, ApJ, 608, L89 Ramirez-Ruiz, E., Merloni, A., & Rees, M. J. 2001, MNRAS, 324, 1147 Ramirez-Ruiz, E., et al. 2005, ApJ, 625, L91
- Resmi, L., et al. 2005, A&A, 440, 477
- Sakamoto, T., et al. 2005, ApJ, 629, 311
- Sazonov, S. Y., Lutovinov, A. A., & Sunyaev, R. A. 2004, Nature, 430, 646
- Scott, M. A., & Readhead, A. C. S. 1977, MNRAS, 180, 539
- Soderberg, A. M., et al. 2004, Nature, 430, 648
- 2006, Nature, 442, 1014
- Stanek, K. Z., et al. 2003, ApJ, 591, L17
- Tagliaferri, G., et al. 2003, IAU Circ. 8308
- Taylor, G. B., Frail, D. A., Berger, E., & Kulkarni, S. R. 2004, ApJ, 609, L1
- Taylor, G. B., et al. 2005, ApJ, 622, 986
- Tiengo, A., et al. 2003, A&A, 409, 983
- -. 2004, A&A, 423, 861
- Tinney, C., et al. 1998, IAU Circ. 6896
- Tominaga, N., et al. 2005, ApJ, 633, L97
- van der Horst, A. J., et al. 2005, ApJ, 634, 1166
- Vanderspek, R., et al., 2004, ApJ, 617, 1251
- Watson, D., et al. 2004, ApJ, 605, L101
- Waxman, E. 2004, ApJ, 605, L97
- Weiler, K. W., et al. 2002, ARA&A, 40, 387
- Woosley, S. E., & Bloom, J. S. 2006, ARA&A, 44, 507
- Woosley, S. E., Eastman, R. G., & Schmidt, B. P. 1999, ApJ, 516, 788
- Zeh, A., Klose, S., & Hartmann, D. H. 2004, ApJ, 609, 952
- Zhang, W., Woosley, S. E., & Heger, A. 2004, ApJ, 608, 365